

Dialogue on Classical and Quantum between mathematician and experimenter

Andrei Khrennikov

International Center for Mathematical Modeling
in Physics and Cognitive Sciences,
MSI, University of Växjö, S-35195, Sweden

Email: Andrei.Khrennikov@msi.vxu.se

Johann Summhammer

Atominstitut der Österreichischen Universitäten
Stadionallee 2, A-1020 Vienna, Austria

E-mail: summhammer@ati.ac.at

February 1, 2008

Abstract

This dialogue took place at Växjö, 19 November 2001. The main aim of our meeting in Växjö was to clarify our viewpoints on foundations of quantum mechanics. The most attractive in our discussion was the extreme difference in our quantum experiences. On one side, pure mathematician (specializing in foundations of probability theory), Andrei Khrennikov; on the other side, pure experimenter (specializing in neutron and electron interferometry), Johann Summhammer. On one hand, an attempt to test mathematical models for larger and larger domains of physical reality. On the other hand, an attempt to create this reality from experimental information - roughly speaking from clicks of detectors.

Andrei: Was the transition from classical to quantum physics a jump-like transition that changed the whole picture of physical reality or just a new application of "classical" statistical physics?

Johann: It was a crucial change of the whole physical ideology. In any case it was not just a new application of classical statistical physics.

Andrei: What are the main distinguishing features of quantum physics?

Johann: I would like to underline two new fundamental features:

(D) Discreteness of quantum observations and finiteness of information that could be obtained via a concrete measurement. In modern language it would be that in finite time only a finite amount of information can be obtained.

(R) Randomness of results of measurements.

Andrei: In fact, I do not see so large difference from classical statistical mechanics. Well, discreteness is the additional condition. But it is natural from the experimentalist viewpoint: a finite precision of each concrete measurement. This problem was deeply investigated already in 19th century: precision, errors and so on. Regarding to (R): I do not see any difference with "classical" statistical physics at all. Already in statistical mechanics we calculate probabilities and averages. It looks that quantum mechanics is just a new domain (it might be very special) of applications of "classical" statistical physics.

Johann: Not at all! First I shall try to explain the role of (D) and then go to (R). Yes, already in 19th century (and even earlier) people paid attention to precision of measurements, errors and so on. However(!), they did not understand the fundamental role of discreteness of information that could be obtained from measurements. Suppose you asked an engineer of 19th century: "Is it possible to measure the length of this stick with arbitrary precision in finite time?" The answer would be: "In principle yes. If we use a very good measurement device, we could do this." So for him the problem was just of a technical matter: we need to improve technology, produce a good device and perform the precise measurement.

Andrei: I am not sure that an engineer of 19th century would behave in such a way. First of all as an engineer he should speak about a measurement with an arbitrary precision, instead of a precise measurement. Moreover, I think that he even would not claim that there exists a kind of "real" length of a stick. On the other hand, it seems that you are right regarding to mathematicians and physicists.

In 19th century they were under great influence of Newtonian model of reality that was based on the use of "continuous quantities" (represented by real numbers in a mathematical model). So it might be that discreteness was surprising for people working with real model of space-time. From this point of view the (D)-argument is just an argument against the use of the mathematical model of physical reality based on real numbers. However, quantum theory did not proceed in this way: the Newtonian "continuous" space-time was transferred automatically into quantum formalism. So the (D)-argument was not so intriguing for experimenters and engineers; it was surprising for mathematicians and physicists working in the continuous Newtonian model. However, in the latter case the (D)-argument did not induce a revolution against the use of continuous quantities in physics - real numbers. It just modifies the formalism (by using matrices) to combine discreteness of results of measurements with "continuous" reality.

Johann: Well, I take this already as an important lesson. Observation gives discrete and finite results, but the reality which we construe from it is left continuous. This is a hint that the reality we can speak about, is the one construed from observations. It is a picture. I think in classical physics we were not forced so rigorously to distinguish between observed data and what we envision from them. Somehow we could think that the data "are" the world.

Now - randomness. There is a crucial difference between randomness in classical statistical physics and quantum physics. Classical randomness is the result of our non sufficient knowledge of objective properties of physical systems. In classical physics you were permitted to think that each classical particle has a definite position at each instant of time. In principle (at least by improving technology) we could find the precise position of a particle. We could not do this for a quantum particle, without strongly influencing its future behaviour. This implies that, the image of a particle as a small something that moves through space, cannot be upheld.

Andrei: So we could not use the classical probability theory that was used in "classical" statistical physics? Just to consider a large ensemble of quantum particles, perform statistical measurement of their properties and find corresponding probabilities and averages?

Johann: No, we could not do this! Classical randomness is, in principle, reducible to deterministic objective realistic description. Quantum stochasticity has a fundamental irreducible character.

Andrei: So classical Ensemble Probability could not work for en-

sembles of quantum systems?

Johann: No it would not. You need to work with the complex probability amplitudes of quantum theory. This has been established in very many experiments. The very working of semiconductor gadgets, to take just one example, is a consequence of electrons following quantum probability rules, not classical rules. [Or take everyday chemistry, like paints and detergents. If you want to understand why they work as they do, you will not be able to do this with classical probability applied to electrons and ions. This is sometimes overlooked when people introduce hidden variable models.

Andrei: What is the origin of such a difference in random behaviour of classical and quantum particles? Do you think (as many) that there is Something very special in quantum systems that produces special quantum probability?

Johann: Yes, I think so.

Andrei: What is this speciality of quantum particles? It may be their wave features?

Johann: I am hesitant to use pictures like wave or particle. Although I do it all the time in the laboratory and when explaining to students, one must be aware that notions of wave, particle and the like are naive projections of our very muscular human experiences onto the world. Although I have recorded "waves" in the form of sinusoidal interference patterns and seen "particles" as flashing spots on a screen, I think it is wrong to attribute "existence" to waves or particles. Quantum physics forces us to recognize that the world simply does not have a "mode of being" such and such. The question what something "really" is, is ill-formulated.

Andrei: I also think that the questions of existence and reality are very important in this discussion. However, I am not sure that these problems are only related to quantum systems. Existence of classical, microscopic, objects is not ABSOLUTE. It seems that the table in this room "exists" only for a special class of observers (including human beings). These observers "create" the table by using their special measurement devices. An observer of the size of an electron or the universe would never "create" such "an element of reality", table.

Johann: From a practical point of view you may be right. But I would argue that any observer, equipped with similar reasoning powers as you and I, could find the same invariants and thus the same structure, when given the observational data which you and I have, and from which we conclude that the particular set of invariants is best

subsumed under the notion "massive impenetrable table". (Incidentally your data and mine are completely different and yet we believe to be talking about the same table.)

Andrei: I think that the crucial role in creation of elements of reality for a special class of observers plays the relative magnitude of perturbations that are produced by measurement devices. I still have the idea that there is not such a large difference between "classical" statistical physics and quantum physics. These two classes of measurements are characterized by different magnitudes of perturbations produced by measurement devices. "Quantum" systems are essentially more sensitive to perturbations of our macro-devices than "classical" systems. Do you agree with such a viewpoint? Negligibly small perturbations - classical physics, relatively large - quantum. It might be that there exist some classes of perturbations that are described neither by quantum nor classical physics.

Johann: No, I cannot follow you here. First, I am uneasy with the term 'perturbation'. It creates the classical image of one object hitting another one, and this is a very muscular view. C.F.von Weizsaecker once said "Hitting and tossing are not primitive notions of physics." I agree with him. I would even say that terms like 'scattering' or 'interaction' will meet with scrutiny in the future. They are also remnants of classical muscular physical thinking. They guide the psychology, which sets the frame for our thinking, in the wrong direction. Aside from that, I think physics can permit only one fundamental description of phenomena. Today this is quantum theory. At a coarser level you will, of course, recover the laws of classical physics. But this will always be understandable as a washing out of interference effects, for instance, when a detector integrates over several interference fringes because they are so narrow, or as uncontrollable interaction and thus entanglement with the greater environment, so that classical probability rules are sufficient to describe what you observe. For practical purposes it may be justified to distinguish between microworld - macroworld. But one cannot see this as a limited applicability of quantum theory.

Andrei: Well, it seems that you as well as I identified classical physics (Newtonian or Maxwellian) physics and classical randomness (ENSEMBLE randomness). I agree with you that we could not use Newtonian picture of physical reality. I even do not claim that quantum particles are really particles, i.e., objects localized in the Euclidean or Minkowski space. These are some objects, "somethings".

The only thing that I defend is that there is no statistical difference between measurements over ensembles of "classical" and "quantum" somethings. These are collections of somethings that have some internal (well, objective) properties. We perform measurements of these properties. However, our measurements produce perturbations. This is the origin of randomness, both classical and quantum? Of course, quantum and classical measurements are characterized by different magnitudes of perturbations. But it is not a consequence of an exotic quantum features. It is just a question of scaling. An observer who has the size of the universe would produce perturbations of quantum magnitude by performing a measurement over an ensemble of our tables.

Johann: No, the different statistical behavior of 'quantum objects', if you permit this term, cannot be reproduced by classical behavior, e.g., when a particle is in a superposition of two different energy states, unless you are willing to introduce an exotic entity like Bohm's quantum potential, which violates special relativity left and right. Moreover, as I just said, I do not like your notion of perturbations. Newtonian physics could not be used in a quantum framework, and notions like forces and perturbations seem to me to be a hindrance. Unless one employs them in the subtle manner as Feynman did: You can think classically, follow each particle through all its conceivable paths, but assign each path a probability AMPLITUDE, not a probability. The immediate question -Why the probability AMPLITUDE? - has no satisfactory explanation until this day, except for symmetry arguments.

Andrei: Are you against my picture of classical+quantum randomness as perturbation-randomness?

Johann: At least your notion of perturbation is not well defined.

Andrei. I agree with you that I have to be more precise with perturbations. I do not consider perturbations as force-like Newtonian perturbations. I am not interested in underlining dynamical model. I speak about Statistical perturbations - perturbations of probability distributions. There are some parameters that determine experimental framework, we can say *context* (the system of physical conditions). It is natural that by varying these parameters we should change context. Physical systems (if you like somethings) would react in a new way to context corresponding to perturbed parameters. We obtain a perturbation of probability distribution. Such a perturbation could be defined in mathematical terms, see [1]. Suppose that small variations

of contex-parameters produce negligible perturbations of probabilistic distributions. We call such experiments - classical. Suppose that small variations of contex-parameters produce non-negligible perturbations of probabilistic distributions. Such a situation we have (in particular, see [1]) in quantum experiments.

Johann: My first reaction would be 'No'. On second thought it seems to me that you are aiming at a new language of speaking about the quantum mechanical time evolution of a system. Not as a continuous transformation of a distribution of probability amplitudes, but as a transformation of probabilities. Isn't the Bohmian picture just that? I am doubtful of the advantage of the distinction 'small perturbation = classical probability' and 'large perturbation = quantum probability'. If you think it through for actual situations, e.g. the double slit experiment with objects of very small and of very large de Broglie wavelength, you may obtain an understanding of what you mean by 'perturbation'. This could be interesting. Although I am sceptical that a deeper explanation of 'why the quantum probability rule?' can be achieved, I am ready to learn. But how about the superposition of alternatives as in the EPR-experiment or the GHZ-experiment? Would perturbation be scaled by how much random spin flip there is? Then you would have the opposite of what you want: No perturbation = quantum probability, large perturbation = classical probability.

[Andrei: The EPR experiment (as well as GHZ are, in fact, experiments on combination of statistical data obtained for distinct complexes of physical conditions, contexts. EPR-contexts are determined by different settings of Stern-Gerlach devices. Here variation of context-parameters, namely parameters of Stern-Gerlach devices, produces non-negligible perturbations. In fact, Bell's inequality is a kind of constraint restricting statistical variations due to variations of context parameters, see [1].

Johann: OK, I see what you mean. The term 'perturbation' tends to lead me astray. It conjures up a 'hitting-tossing' image.

Andrei. Recently I read Einstein's papers on blackbody radiation and photoelectric effect. First of all I was surprised by clearness of presentation of ideas (especially, compare to Bohr, Heisenberg, Dirac). So Einstein derived "quantum probabilistic distributions" by using methods of classical statistical physics. These were the standard statistical manipulations with ensembles of particles + discreteness. However, the (D)-argument does not look so non classical. OK, discrete values, but ordinary probability?

Johann: In hindsight I would say that Einstein got the right result, because superposition of different number states of the electromagnetic field did not play a role in the problem. If it had, as for instance for a field in a cavity consisting of a superposition of a 4-photon and a 5-photon state and the question is what phase shift will an atom pick up when it passes through, he would probably have insisted that there are either really 4 photons or really 5 photons in the cavity, if I apply the reasoning he used in the EPR-paper. This would have led to the wrong phase shift for the atom.

Andrei: Do you think that Einstein obtained "quantum probabilistic distributions" just accidentally? Did he miss the superposition features of quantum systems just by accident?

Johann: Yes, it seems so...

Andrei: I think that you would agree that in each particular quantum experiment, as a measurement of some fixed physical observable, we get classical statistical data and we operate with this data in the same way as with the statistical data obtained in classical experiments. It seems that the crucial probabilistic difference between classical and quantum is demonstrated in the interference experiments. Let us consider the two slit experiment. As I understood, we both do not try to reduce the problem to wave description. For you: a particle passes just one slit (when both are open) or in some way both slits?

Johann: First, I agree, that any quantum experiment of fixed conditions gives data which follow a well known classical probability distribution: binomial, multinomial, Poissonian. The problem of understanding arises when you try to make sense of several such experiments in which one or several parameters have been changed, and you attempt a classical visualization of 'what the particle(s) did'. As to the double slit experiment, it is clear that the particle picks up information (=transformation of probability amplitude) along BOTH paths and this determines the probability where it will end up on the screen. But again, I am doubtful of this imagery. Because, any attempt of verifying where the particle went changes the experimental condition to such a degree, that interference vanishes (here you have a case of 'strong perturbation = classical probability'). The information, where the particle went, must be principally nonexistent in the universe, then you get (maximum) interference. Now, if this information is nowhere available, what sense does it make to stubbornly hold on to the picture 'but it must have gone somewhere'? I take this lesson to mean that the old and taken-for-granted view of the everyday

world, which we think to be well defined and always there, is wrong and untenable. That, what is there, is of a lot more abstract nature. Personally, I tend towards information as the new key concept. But not information about 'things'. Information about possible values of properties, and sets of properties, which are not there 'by themselves', but which WE define on the basis of symmetries. Which is almost as good as saying they are there 'by themselves', because the kind of symmetries we can think of to categorize observations is not our free invention, but is determined by the laws of thought. And these are definitely not our own creation, but are structures which we run up against in our minds.

Andrei: I am not against reality as information reality. We can consider quantum as well as classical objects as just collections of information. There exist objective laws of evolution and interaction of information; in particular, the laws of thought.

Johann: I would not use a mechanical idea like 'information interaction', otherwise I agree.

Andrei: Well, we turn back to probability. I consider interference (classical as well as quantum) as "interference" of two contexts or two alternatives. The two slit experiment for one fixed experimental setting, i.e. both slits are open, could not demonstrate special "quantum features." Interference fringes by themselves are not exhibitions of special quantum nature. For instance, we can get interference patterns in the following classical variant of the two slit experiment.

Charged macroscopic bodies move through charged screen having two slits. We get a complex interference like distribution of bodies on the registration screen. What is the root of this interference? The special character of interaction between charged bodies and charged surface of the first screen. Of course, in quantum two slit experiment we do not have such internal (Newtonian) description of interaction between a particle and the screen; at least in the conventional quantum formalism. There is no quantum charge... However, we could use, for example, Bohmian mechanics (pilot wave formalism) with its quantum potential or quantum field given by the wave function.

Johann: Yes, you can get such periodic-looking classical particle scattering patterns in certain classical experiments. But this has nothing to do with quantum interference. As to Bohmian mechanics, I view this as a perhaps desperate attempt to save our classical intuitive pictures. What I criticize is that it misses the message of quantum theory, which is similar to the message of special and general relativ-

ity theory. Namely that, our classical intuition, in which the world is made up of distinct things, which exercise forces on each other, and all this happens in a well-set three-dimensional expanse, is simply wrong. I read the message nature gives us in these theories as: "Thou shalt not make an image of mine!", a warning, which is probably already in the Old Testament. If we teach students the Bohmian mechanics, we may help them in the first few steps into quantum theory, but the damage we do by distracting them from wider schemes of thought is in the long run much greater. Today there exist computer programs which let you view the temporal evolution of the probability density obtained from the Schrödinger equation for one or several particles for many different Hamiltonians. This can help to form a new kind of intuition.

Andrei: In fact, I do not claim the Bohmian mechanics provides the right internal description of quantum phenomena. But at least this is one of possible models.

Johann: Models are certainly needed to do practical research. But I would emphasize seeking in data the symmetries, sets of symmetries, sets of sets of symmetries, and so on.

Andrei: We now turn back to the quantum two slit experiment. So interference by itself is not the crucial exhibition of QUANTUM. Troubles with classical picture, in fact, classical probability, start when we play with a few experimental settings (contexts) in the two slit experiment: (C_{12}) both slits open; (C_1) only the first slit is open; (C_2) only the second slit is open. If we add probabilities obtained for contexts (C_1) and (C_2) we do not get the probability for the context (C_{12}) . Classical rule for the addition of probabilities of two alternatives (first or second slit):

$$P = P_1 + P_2. \quad (1)$$

is violated. In experiments we get so called quantum rule for addition of probabilities:

$$P = P_1 + P_2 + 2\sqrt{P_1 P_2} \cos \theta. \quad (2)$$

Many people consider this fact as the real trouble with classical probability. They start to think that there is something wrong with the image of electron passing through both slits. Moreover, this is the root of the wave approach to quantum mechanics.

However, we will be in trouble only if we forget about perturbation of probabilistic distributions induced by the transition from one

context to others. In fact, there are three different contexts that are involved in the consideration: (C_{12}) and (C_1) , (C_2) . "Quantum systems" are sensitive to variations of context parameters; in our case: opening and closing of slits. For "quantum systems" the complex of physical conditions (C_{12}) differs strongly from complexes (C_1) and (C_2) . In fact, (2) is just the transformation that connects probability distributions related to different contexts. The classical rule (1) was derived for the fixed context - in conventional axiomatics of probability theory, Kolmogorov, 1933. Thus if we control context dependence in the right way, then we shall not obtain any contradiction between quantum statistical data and classical Kolmogorov probability theory. The main problem in the right probabilistic understanding of quantum statistical experiments is that people try to apply theory of probability that was developed for one fixed context to statistical data obtain in a few different contexts.

Johann: When you speak of contexts rather than perturbation, I can follow you more easily. Because then you sound like Niels Bohr, who insisted that the whole experimental arrangement determines the probability distribution for the possible outcomes. He is absolutely right, if you consider that the slightest change in the experiment must be entered as a corresponding change in the Hamiltonian which goes into the Schrödinger equation.

As to the Kolmogorovian axioms of probability theory, you seem to imply that quantum theory does not contradict them. I agree with you. The particle was sent onto a diaphragm with two slits and later ended up at some point on the screen. This is only one event, observed under one context, not an event that can have happened under two different contexts. This is not possible. Any given experimental situation is always only one context. Nevertheless, the task remains, how to explain the existence of a lawful relation between the probabilities observed in the three different contexts. (And jokingly one might ask, why should two contexts suffice to tell us something about a third? Perhaps it takes three, to fully predict the outcomes under a fourth? Then we would need a three-slit experiment to capture the whole mystery of quantum physics.)

I have a way of viewing this, which is based on the idea that, more observations shall permit more accurate predictions. This is a natural assumption for an experimenter. But it leads to strong constraints on how the probabilities observed under two different contexts can functionally be related to the probabilities found under a third one.

In the two-slit case it means that, if you measure the probabilities for the particle to arrive at a particular spot on the screen in two different contexts, you should be able to predict the probability, or at least a function of it, under the third context ever more accurately, the more trials you have done to measure the first two. If you follow this through, you get a real and a complex way of describing this, and the latter is the quantum theoretical rule. But you do not get the classical rule [2]. For this reason I am interested in questions like "Which theory permits the most accurate predictions from the information that is factually available now?" as an entry point to understanding quantum theory.

Maybe I can comment a little more on the relation between classical and quantum probabilistic behavior. I think at bottom there is only quantum probability. Classical probabilistic behavior appears because one does not look closely, and so collects 'dirty' facts, either by not controlling the context sufficiently, so that the observed data are actually collected under many different contexts, or by resolving the measurement results with insufficient fine grading. Let us take the double slit experiment again. If you do this experiment with an ensemble of particles which do not all have the same energy, but a wide spread of energies, the probability distribution on the screen will look very much as if you had used classical particles of the same wide energy distribution. So you have actually averaged over many different contexts, because every specific energy requires a different preparation context. On the other hand, insufficient fine grading of the measurement would mean, that you have interference fringes on the screen, which are narrower than the grains of the emulsion of the photographic plate, or the size of the individual pixels, if you use a modern digital array chip as detection screen. I have no doubt that all phenomena observed in classical probabilistic physics can be traced back to 'sloppy' context definition and 'sloppy' data registration, and mostly to a combination of both. And the probability rule $P_1 + P_2 = P_{12}$ works there only, because the sloppiness of experimentation is compensated with a really wrong application of the notion of 'event', as it appears in Kolmogorov's third axiom. Of course, in classical physics both mistakes were done unknowingly. I consider it an accident that the classical rule works in the everyday world, rather than a self-evident truth.

Andrei: It seems to me that in fact the discovery of quantum formalism was merely a discovery in probability theory and not really in physics. Quantum formalism, at least Hilbert space calculus, is

a new mathematical theory to work with statistical data belonging to a few different contexts. Moreover, it was discovered that non-linear probabilistic transformation (2) can be represented as a linear transformation (for square roots of probabilities) in a complex Hilbert space. Typically the main attention is paid to this Hilbert space calculus. But I think that the crucial point was the derivation (at the beginning purely experimental) of transformation (2) connecting probabilities with respect to three different contexts. In fact, linear algebra can be easily derived from this transformation. Everybody familiar with the elementary geometry will see that (2) just the well known cos-theorem. This is the rule to find the third side in a triangle if we know lengths of two other sides and the angle θ between them:

$$c^2 = a^2 + b^2 - 2ab \cos \theta .$$

or if we want to have "+" before $\cos$ we use so called *parallelogram law*:

$$c^2 = a^2 + b^2 + 2ab \cos \theta . \quad (3)$$

Here c is the diagonal of the parallelogram with sides a and b and the angle θ between these sides. Of course, the parallelogram law is just the law of linear (two dimensional Hilbert space) algebra: for finding the length c of the sum $\mathbf{c}$ of vectors $\mathbf{a}$ and $\mathbf{b}$ having lengths a and b and the angle θ between them.

Johann: I would disagree with the statement that the quantum formalism was not a discovery in physics. As I said before, it implied a big change in our approach to understanding the world and we have not yet fully explored all consequences. But I can agree with you on Hilbert space as a new tool for dealing with statistical data.

References.

[1] A. Khrennikov, Linear representations of probabilistic transformations induced by context transitions. *J. Phys. A: Math. Gen.*, **34**, 1-17 (2001).

[2] J. Summhammer, Maximum predictive power and the superposition principle. *Int. J. Theor. Phys.*, **33**, 171 (1994).